

# Joint effect of high blood pressure and physical inactive on diabetes mellitus: a population-based cross-sectional survey

*anonymous marking enabled*

---

**Submission date:** 04-Jun-2022 03:27PM (UTC+0700)

**Submission ID:** 1850225612

**File name:** 2021\_Joint\_effect\_of\_high\_blood\_pressure.pdf (149.71K)

**Word count:** 5058

**Character count:** 25824



# Joint effect of high blood pressure and physical inactive on diabetes mellitus: a population-based cross-sectional survey

2

NIKSON SITORUS<sup>1</sup>, APRIZAL SATRIA HANAFI<sup>2</sup>, DEMSA SIMBOLON<sup>3</sup>, MARTINA<sup>2</sup>

<sup>1</sup> Center for Research and Development of Public Health Efforts, National Institute of Health Research and Development, Ministry of Health RI, Jakarta, Indonesia; <sup>2</sup> Alumnus of Master of Epidemiology Student, School of Public Health, University of Indonesia, Depok, Indonesia; <sup>3</sup> Department of Public Health, Health Polytechnic of Bengkulu Ministry of Health, Bengkulu, Indonesia

## Keywords

Joint effect • Blood pressure • Physical inactivity • Diabetes mellitus • Indonesia

## Summary

**Introduction.** The relationship of high blood pressure and physical inactivity to diabetes mellitus is well known, but not many studies have known the joint effect of the two in causing diabetes mellitus. This study aims to evaluate the joint effect of high blood pressure and less physical activity against Diabetes Mellitus (DM) in Indonesia.

**Methods.** This is a cross-sectional study. Subjects in this study were the age group  $\geq 21$  years old who were followed by the interview. We investigated factors related to DM in Indonesia associated with blood pressure and physical activity by controlling other confounding variables. Statistical analyses were conducted using logistic regression. Age, sex, education level, marital status, occu-

pation, body mass index, residence area, stress, fruit, and vegetable consumption were adjusted for in the multivariate model.

**Results.** The prevalence of DM was 3.86% among respondents. Multivariate analysis showed that people who had hypertension and less physical activity had a risk of 3.68 (95% CI, 2.43-5.34) times having DM. People who had hypertension and enough physical activity had a risk of 2.33 (95% CI, 1.65-6.43) times having DM. While people who do not have hypertension and had less physical activity had a risk of 1.81 (95% CI, 1.34-3.62) times.

**Conclusions.** People with hypertension and less physical activity have the greatest risk of developing DM.

## Introduction

In 2014, according to WHO there were 422 million adults aged over 18 who lived with Diabetes Mellitus (DM) [1]. Prevalence of DM in Indonesia has continued to increase, from 5.7% in 2007, to 6.9% in 2013, and increased again to 8.5% in 2018 [2-4]. Indonesia is the 4<sup>th</sup> country with the highest prevalence of DM in the world. Even the number of people with DM continues to increase from year to year. WHO data estimates that the number of people with DM in Indonesia will increase significantly to 21.3 million in the next 2030 [5].

Lifestyle factors and clinical factors are among the other factors that have a major influence on the incidence of DM. According to the study from Coldberg (2016) and Williams (2013) high blood pressure and less physical activity were the main predictor factors that trigger an increase in blood sugar levels. So, the two factors must be controlled as prevention efforts [6].

Based on these data it can be seen that the prevalence of DM patients increases every year in Indonesia and the joint effects of blood pressure and physical activity in influencing DM events have never been done. In previous study by Hanafi and Prihartono (2018), a similar study was carried out with different study outcomes [7]. This study aims to find the joint effects of blood pressure and physical activity with DM by controlling other variables

such as age, sex, marital status, education level, occupation, residence area, body mass index, stress, vegetable consumption, and fruit consumption.

## Methods

### ETHICAL CONSIDERATIONS

The IFLS-5 survey procedures had been approved by Institutional Review Boards (IRBs) in the United States at Rand Corporation, Santa Monica, California and in Indonesia at Ethics Committees of Gadjah Mada University.

### STUDY DESIGN

This study uses a cross-sectional design using data from the Indonesian Family Life Survey-5 [8]. The survey collected information on individual, household and community level data using multistage stratified random sampling. IFLS is a longitudinal household survey involving 7<sup>th</sup> questionnaire and anthropometric measurements, and which was collected under the supervision of the Rand Corporation. IFLS-5 was conducted in 13 provinces in Indonesia [9].

IFLS-5 was conducted in September 2014-March 2015 on 50,148 individuals. The study population was the population who became the subject of IFLS-5 research

in 2014. While the sample was the age group  $\geq 21$  years who followed the interview and had questionnaire data on important variables [10].

### STUDY VARIABLE

We include demographic information, individual characteristics and behavioral factors as confounding. We categorize the level of education completed by respondents to low (under Senior High School), middle (Senior High School) and high (College or University), while marital status was classified as single, married, separated, live divorced, death divorced. Occupations were categorized as working and not working.

Physical activity was assessed through a series of questions a brief form modified from the International Physical Activity Questionnaire (IPAQ) on the type and time of physical activity involved in, in all parts of life: work, home and exercise and then classified as enough and less physical activity [11].

DM is assessed through questions ever diagnosed or not done by doctors or paramedics. We also measured respondents' fiber consumption in the past week, which was seen from the consumption of fruits and vegetables. Body mass index (BMI)  $< 27 \text{ kg/m}^2$ : normal weight; and  $\geq 27.0 \text{ kg/m}^2$ : obesity derived from the height and weight measured during the physical examination, these criteria were determined based on the Ministry of Health of the Republic of Indonesia in 2013. Height measured by the Seca plastic height board model 213 and weight was measured using Camry model EB 1003 scale. In this study the measurement of body weight and height was carried out by the interviewer or enumerator who was competent in their field and had received previous training.


Blood pressure was measured 3 times at an individual, using Omron meter HEM 7203. The first measurement was done at the beginning of the interview with the next

two steps taken during the interview. The average of the 3 measurements was used for the current analysis. According to the JNC 7 blood pressure was categorized into 4 levels, namely normal ( $< 120/80 \text{ mmHg}$ ), pre-hypertension ( $120\text{-}139/80\text{-}89 \text{ mmHg}$ ), hypertension stage 1 ( $140\text{-}159/90\text{-}99 \text{ mmHg}$ ), and hypertension stage 2 ( $\geq 160/100 \text{ mmHg}$ ). We classify respondents as hypertension if their blood pressure  $\geq 140/90 \text{ mmHg}$  based on the criteria of JNC 7. Blood pressure measurement was carried out by the interviewer or enumerator who was competent in their field. Only respondents with complete information and blood pressure measurements were taken 3 times included in the analysis. After processing the data all of our study variables continued by making the joint variable of blood pressure and physical activity into one variable. The joint variables are divided into 4 categories, namely groups of people who are not hypertensive and have enough physical activity, groups of people who are not hypertensive and have less physical activity, groups of people who are hypertensive and have enough physical activity, and groups of people who are hypertensive and have less physical activity.

### STATISTICAL ANALYSIS

Logistic regression was performed to calculate the risk in all age groups. This study includes age, sex, education level, marital status, occupation, body mass index, residence area, stress, fruit and vegetable consumption and as potential confounders variables by including them in multivariable analysis between blood pressure and physical activity to DM. If there is a difference of more than 10% between POR crude and POR adjusted then these variables were considered as confounding variables and not included in the next model. The same procedure was used to estimate adjusted odds ratio (and 95% CI) for DM [12]. Finally, the joint effect (and 95% CI) of hyper-

Fig 1. Selection of study sample flowchart.



tension and physical activity, individual effect of hypertension among people with enough physical activity, and effect of people with less physical activity among non hypertension on DM were evaluated (Fig. 1).

**Results**

The description of each study variable can be seen in Table I. Of the 14,053 respondents, the proportion of DM in Indonesia was 3.86%. While the proportion of hypertension and less physical activity was 18.50% and 21.20%, respectively.

Table I shows that the majority of respondents were 21-44 years old (62.61%), women (54.52%), married (78.70%), low education (85.76%), working (81.80%), living in rural areas (54.79%), not hypertensive (81.50%), not obese (73.81%), enough physical activity (78.80%), not stressed (62.72%), consuming vegetables 7 days/week (48.39%), and consuming fruits 7 days / weeks (39.86%). The results of joint variables of blood pressure and physical activity showed that most respondents were in the category of non-hypertensive and enough physical activity (64.93%) and the least in the hypertension and less physical activity group (4.63%).

Based on Table II shows that the proportion of DM is highest in the 45-59 year age group (51.48%), women (55.54%), married people (83.76%), low education (74.17 %), people who live in urban areas (67.16%), obese people (54.61%), people who have enough physical activity (62.18%). Variables of age, sex, education level, marital status, residence area, blood pressure, body mass index, physical activity, fruit and vegetable consumption were significantly associated with DM with p value < 0.05. While the occupation and stress variables do not show a significant relationship with a p value > 0.05.

Based on the joint variable blood pressure and physical activity the proportion of the highest diabetes mellitus is indeed in the group of people who are not hypertension and have enough physical activity (33.95%). However, this is due to the fact that the proportion in this group is the highest, namely 64.93% (Tab. II). Interestingly, the group with the second and third highest proportion of DM was a group of people with hypertension and enough physical activity (28.23%) and groups of people with hypertension and less physical activity (19.93%). While the group of people without hypertension and less activity the least proportion of DM. So, it can be concluded that hypertension is a significant factor in influencing the proportion of DM than physical activity (Tab. III).

Table IV shows that the highest risk of DM is in the group of people who have hypertension and less physical activity which is 3.68 times, while those in hypertension and less physical activity risky 2.33 times, in groups of people who are not hypertension and have less physical activity risky 1.81 times greater with a group of people who are not hypertension and have enough physical activity as a reference.

Tab. I. Respondents features.

| Characteristic  | Total  | Percentage |
|---|--------|------------|
| <b>Diabetes mellitus</b>                                      |        |            |
| Yes   | 542    | 3.86       |
| No  | 13,511 | 96.14      |
| <b>Age (years)</b>  |        |            |
| 21-44 (adults)  | 8,799  | 62.61      |
| 45-59 (middle)  | 3,543  | 25.21      |
| 60-74 (elderly)   | 1,498  | 10.66      |
| 75-90 (old)   | 213    | 1.52       |
| <b>Sex</b>  |        |            |
| Male  | 6,392  | 45.48      |
| Female  | 7,661  | 54.52      |
| <b>Marital status</b>   |        |            |
| Single  | 1,598  | 11.37      |
| Married   | 11,060 | 78.70      |
| Separated   | 80     | 0.57       |
| Live divorced   | 297    | 2.11       |
| Death divorced  | 1,018  | 7.24       |
| <b>Education</b>  |        |            |
| High  | 1,503  | 10.70      |
| Middle  | 498    | 3.54       |
| Low   | 12,052 | 85.76      |
| <b>Occupation</b>   |        |            |
| Yes   | 11,496 | 81.80      |
| No  | 2,557  | 18.20      |
| <b>Residence area</b>   |        |            |
| Rural   | 7,700  | 54.79      |
| Urban   | 6,353  | 45.21      |
| <b>Blood pressure</b>   |        |            |
| Non-hypertension  | 11,453 | 81.50      |
| Hypertension  | 2,600  | 18.50      |
| <b>Body mass index</b>  |        |            |
| Normal  | 10,372 | 73.81      |
| Obesity   | 3,681  | 26.19      |
| <b>Physical activity</b>                                      |        |            |
| Enough  | 11,074 | 78.80      |
| Less  | 2,979  | 21.20      |
| <b>Stress</b>   |        |            |
| No  | 8,814  | 62.72      |
| Yes   | 5,239  | 37.28      |
| <b>Vegetable consumption</b>                                  |        |            |
| 7/week  | 6,800  | 48.39      |
| 4-6/week  | 3,725  | 26.51      |
| 1-3/week  | 3,058  | 21.76      |
| Never   | 470    | 3.34       |
| <b>Fruit consumption</b>                                      |        |            |
| 7/week  | 5,601  | 39.86      |
| 4-6/week  | 3,919  | 27.89      |
| 1-3/week  | 3,464  | 24.65      |
| Never   | 1,069  | 7.61       |
| <b>Joint variable of blood pressure and physical activity</b> |        |            |
| Non-hypertension + enough                                     | 9,125  | 64.93      |
| Non-hypertension + less                                       | 2,328  | 16.57      |
| Hypertension + enough   | 1,949  | 13.87      |
| Hypertension + less   | 651    | 4.63       |

Tab. II. Frequency of diabetes mellitus according to individual characteristics.

| Characteristics              | Diabetes mellitus |       | Non-diabetes mellitus |       | P value | POR  | 95% CI    |
|------------------------------|-------------------|-------|-----------------------|-------|---------|------|-----------|
|                              | N = 542           | %     | N = 13,512            | %     |         |      |           |
| <b>Age (years)</b>           |                   |       |                       |       |         |      |           |
| 21-44 (adults)               | 180               | 33.21 | 8,619                 | 63.79 |         | 1    | 1         |
| 45-59 (middle)               | 279               | 51.48 | 3,264                 | 24.16 | < 0.001 | 1.68 | 1.21-2.15 |
| 60-74 (elderly)              | 80                | 14.76 | 1,418                 | 10.50 | < 0.001 | 2.70 | 2.06-3.53 |
| 75-90 (old)                  | 3                 | 0.55  | 210                   | 1.55  | < 0.001 | 4.09 | 3.37-4.95 |
| <b>Sex</b>                   |                   |       |                       |       |         |      |           |
| Male                         | 241               | 44.46 | 6,151                 | 45.53 |         | 1    | 1         |
| Female                       | 301               | 55.54 | 7,360                 | 54.47 | < 0.001 | 1.04 | 0.87-1.24 |
| <b>Marital status</b>        |                   |       |                       |       |         |      |           |
| Single                       | 17                | 3.14  | 1,581                 | 11.70 |         | 1    | 1         |
| Married                      | 454               | 83.76 | 10,606                | 78.50 | < 0.001 | 3.98 | 2.44-6.47 |
| Separated                    | 1                 | 0.18  | 79                    | 0.58  | 0.011   | 1.17 | 1.05-8.95 |
| Live divorced                | 14                | 2.58  | 283                   | 2.09  | 0.045   | 4.60 | 2.24-9.43 |
| Death divorced               | 56                | 10.33 | 962                   | 7.12  | 0.005   | 5.41 | 3.12-9.37 |
| <b>Education</b>             |                   |       |                       |       |         |      |           |
| High                         | 98                | 18.08 | 1,405                 | 10.40 |         | 1    | 1         |
| Middle                       | 42                | 7.75  | 456                   | 3.38  | 0.035   | 2.02 | 1.60-2.53 |
| Low                          | 402               | 74.17 | 11,650                | 86.23 | 0.004   | 2.66 | 1.91-3.71 |
| <b>Occupation</b>            |                   |       |                       |       |         |      |           |
| Yes                          | 11,135            | 82.41 | 361                   | 66.61 |         | 1    | 1         |
| No                           | 2,376             | 17.59 | 181                   | 33.39 | 0.051   | 2.34 | 1.95-2.82 |
| <b>Residence area</b>        |                   |       |                       |       |         |      |           |
| Rural                        | 178               | 32.84 | 7,522                 | 55.67 |         | 1    | 1         |
| Urban                        | 364               | 67.16 | 5,989                 | 44.33 | < 0.001 | 2.56 | 2.14-3.08 |
| <b>Blood pressure</b>        |                   |       |                       |       |         |      |           |
| Non-hypertensive             | 281               | 51.85 | 11,172                | 82.69 |         | 1    | 1         |
| Hypertensive                 | 261               | 48.15 | 2,339                 | 17.31 | < 0.001 | 2.22 | 1.18-3.26 |
| <b>Body mass index</b>       |                   |       |                       |       |         |      |           |
| Normal                       | 246               | 45.39 | 10,126                | 74.95 |         | 1    | 1         |
| Obesity                      | 296               | 54.61 | 3,385                 | 25.05 | < 0.001 | 2.31 | 1.46-3.67 |
| <b>Physical activity</b>     |                   |       |                       |       |         |      |           |
| Enough                       | 337               | 62.18 | 10,373                | 79.47 |         | 1    | 1         |
| Less                         | 205               | 37.82 | 2,774                 | 20.53 | < 0.001 | 1.76 | 1.01-3.06 |
| <b>Stress</b>                |                   |       |                       |       |         |      |           |
| No                           | 322               | 59.41 | 8,492                 | 62.85 |         | 1    | 1         |
| Yes                          | 220               | 40.49 | 5,019                 | 37.15 | 0.066   | 1.15 | 1.07-1.37 |
| <b>Vegetable consumption</b> |                   |       |                       |       |         |      |           |
| 7/week                       | 257               | 47.42 | 6,543                 | 48.43 |         | 1    | 1         |
| 4-6/week                     | 154               | 28.41 | 3,571                 | 26.43 | < 0.001 | 1.19 | 0.69-2.04 |
| 1-3/week                     | 116               | 21.40 | 2,942                 | 21.77 | < 0.001 | 1.30 | 1.16-2.28 |
| Never                        | 15                | 2.77  | 455                   | 3.37  | < 0.001 | 2.11 | 1.17-2.86 |
| <b>Fruit consumption</b>     |                   |       |                       |       |         |      |           |
| 7/week                       | 246               | 45.39 | 5,355                 | 39.63 |         | 1    | 1         |
| 4-6/week                     | 158               | 29.15 | 3,761                 | 27.84 | < 0.001 | 1.31 | 0.90-1.93 |
| 1-3/week                     | 105               | 19.37 | 3,359                 | 24.86 | < 0.001 | 1.44 | 1.09-2.12 |
| Never                        | 33                | 6.09  | 1,036                 | 7.67  | < 0.001 | 2.44 | 1.89-3.08 |

Tab. III. Frequency of diabetes mellitus according to joint variable of blood pressure and physical activity.

| Characteristics           | Diabetes mellitus |       | Non-diabetes mellitus |       | POR  | 95% CI    |
|---------------------------|-------------------|-------|-----------------------|-------|------|-----------|
|                           | N = 542           | %     | N = 13,512            | %     |      |           |
| Non-hypertension + enough | 184               | 33.95 | 8,941                 | 66.18 | 1    | 1         |
| Non-hypertension + less   | 97                | 17.90 | 2,231                 | 16.51 | 1.85 | 1.17-3.53 |
| Hypertension + enough     | 153               | 28.23 | 1,796                 | 13.29 | 2.52 | 1.86-5.27 |
| Hypertension + less       | 108               | 19.93 | 543                   | 4.02  | 3.82 | 2.54-9.35 |

Tab. IV. Final model of joint variable of blood pressure and physical activity against diabetes mellitus.

| Joint variable of blood pressure and physical activity | Diabetes mellitus |       | Non-diabetes mellitus |       | POR (95% CI)     |
|--|-------------------|-------|-----------------------|-------|------------------|
|  | N = 542           | %     | N = 13,512            | %     |                  |
| Non-hypertension + enough                              | 184               | 33.95 | 8,941                 | 66.18 | 1.00 (reference) |
| Non-hypertension + less                                | 97                | 17.90 | 2,231                 | 16.51 | 1.81 (1.34-3.62) |
| Hypertension + enough                                  | 153               | 28.23 | 1,796                 | 13.29 | 2.33 (1.65-6.43) |
| Hypertension + less                                    | 108               | 19.93 | 543                   | 4.02  | 3.68 (2.43-5.34) |

Adjusted by age, sex, education, occupation, residence area, body mass index, and fruit and vegetable consumption.

## Discussion

The starting point for healthy living with diabetes is an early diagnosis, the longer a person lives with undiagnosed and untreated diabetes, the worse the health outcome. For those diagnosed with diabetes, a series of interventions can reduce the risk of bad prognosis diabetes, regardless of what type of diabetes they may have. These interventions include blood pressure control, blood glucose, through a combination of diet, physical activity and, if necessary, treatment, to facilitate early control [1].

Our study shows that (48.15%) people with DM have hypertension. Most people with DM were female. This is in line with several other studies that show that women suffer more from DM [13-17]. Most respondents have low education and working [18, 19]. This study also shows that most people with DM are adults (21-44 years) and married. This is in line with several other studies. The results of the same study were also mentioned by other studies [18].

This study shows that in non-diabetes mellitus patients have enough physical activity than less physical activity. Other studies suggest that physical activity can improve blood sugar control [20].

This cross-tabulation analysis also shows that the variables of age, sex, education level, marital status, residence area, blood pressure, body mass index, physical activity, fruit and vegetable consumption are significantly associated with DM. While the occupation and stress variables do not show a significant relationship.

Physical activity includes all movements that increase energy use. Exercise improves blood glucose control in DM, reduces cardiovascular risk factors, contributes to weight loss, and improves well-being [21, 22]. Enough physical activity can prevent or delay the development of diabetes [23]. Regular exercise also has considerable health benefits in people with diabetes (e.g., increased cardiovascular fitness, muscle strength, insulin sensitivity etc. [24]. Challenges related to blood glucose management. Insulin in the muscles and liver can be modified immediately by physical activity and regular physical activity [25]. Aerobic exercise increases muscle glucose up to 5-fold. After exercise, glucose uptake remains increased by insulin-independent (2 hours) and insulin dependent (up to 48 hours) [26].

Physical activity is not the only trigger factor for DM in the equation below explained about the joint effects of blood pressure and physical activity on the occurrence of DM. The pathophysiological mechanism that explains

the relationship between hypertension and the incidence of DM, not yet clear. However high blood pressure has been shown to induce microvascular dysfunction, which can contribute to the pathophysiology of the development of diabetes [27, 28]. Endothelial dysfunction associated with insulin resistance is also associated with hypertension, and biomarkers of endothelial dysfunction are predictors of DM [29].

Elevated blood pressure values are a common finding in patients with DM and are thought to reflect, at least in part, the impact of the underlying insulin resistance on the vasculature and kidney [30]. On the contrary, accumulating evidence suggests that disturbances in carbohydrate metabolism are more common in hypertensive individuals [31, 32]. Thereby indicating that the pathogenic relationship between DM and hypertension is actually bidirectional.

In the multivariate analysis of joint variables of blood pressure and physical activity was found that hypertension had a greater effect of 2.33 times in causing DM than less physical activity ie 1.81. However, the risk of DM increases significantly, which is 3.68 times when hypertension and less physical activity appear together. In the above results, the percentage of the increased risk of DM events can be calculated when hypertension and less physical activity appear together as follows:

- $(3.68-1) = (2.33-1) + (1.81-1)$ ;
- $2.68 = 1.33 + 0.81$ ;
- $2.68 = 2.14$ ;
- $2.68 > 2.14$ ;
- $2.68-2.14 / 2.68 = 20.14\%$ .

This means that the risk for developing DM will increase by 20.14% when hypertension and less physical activity appear simultaneously due to the interaction of both.

This study has limitations, because this is a Cross-Sectional study, so it cannot determine causal relationships. Longitudinal studies are needed to assess the joint effect of blood pressure and physical activity on DM to draw strong conclusions about the causal pathways of this relationship.

## Conclusions

The proportion of DM in Indonesia who became respondents in IFLS-5 is 3.86%. The combination of hypertension and less physical activity have a risk of 3.86 times to suffer from DM compared to those who not hypertension and have enough physical activity. Hypertension and less physical activity together show a greater

association with DM than hypertension or less physical activity alone. The continued increase in DM prevalence makes it necessary to increase health promotion efforts including the addition of nutrition counseling and counseling as well as joint sports activities (gymnastics) in integrated coaching activities. Communities, especially those classified as high-risk (hypertension and less physical activity) can realize the importance of independently performing DM screening in this case was blood pressure, blood glucose level, general obesity of body weight and height.

## Acknowledgements

We are thankful to the Center for Research and Development of Public Health Efforts, National Institute of Health Research and Development, Ministry of Health RI, Jakarta, Indonesia. We are grateful to RAND for providing data.

Funding sources: this study was supported by Center for Research and Development of Public Health Efforts, National Institute of Health Research and Development, Ministry of Health RI, Jakarta, Indonesia.

## Conflict of interest statement

The authors declare no conflict of interest.

## Authors' contributions

NS: conceived of the presented idea, collect the data, performed the analysis. ASH: conceived of the presented idea, conceived and designed the analysis, contributed data or analysis tools, performed the analysis, contributed to the interpretation of the results. DS: verified the analytical methods, designed the model and the computational framework, other contribution. M: developed the theory and performed the computations, performed the analysis, derived the models and analyzed the data, wrote the paper. All authors discussed the results and contributed to the final manuscript.

## References

- [1] WHO. Global report on diabetes, 2014. Available from: [https://apps.who.int/iris/bitstream/handle/10665/204871/9789241565257\\_eng.pdf;jsessionid=58FB19153D48798699A14CEBD67E62B1?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/204871/9789241565257_eng.pdf;jsessionid=58FB19153D48798699A14CEBD67E62B1?sequence=1)
- [2] Ministry of Health RI. Basic Health Research (Riset Kesehatan Dasar). Jakarta: National Institute of Health Research and Development 2007.
- [3] Ministry of Health RI. Basic Health Research (Riset Kesehatan Dasar). Jakarta: National Institute of Health Research and Development 2013.
- [4] Ministry of Health RI. Basic Health Research (Riset Kesehatan Dasar): main result. Jakarta: National Institute of Health Research and Development 2018.
- [5] Siena I. The number of diabetics in Indonesia is astounding.

- Yayasan Sinergi Muda Indonesia 2017. Available from: <https://mudazine.com/ibnusia/penderita-diabetes>
- [6] Williams B. Blood pressure and diabetes: a fatal attraction. *Eur Heart J* 2013;34:3395-7. <https://doi.org/10.1093/eurheartj/ehz365>
  - [7] Hanafi AS and Prihartono NA. Joint effect of obesity and cigarette smoking against hypertension stage 1 among men adults: finding from the Indonesian Family Life Survey-5. *Indian Journal of Public Health Research and Development* 2018;9:673-8. <https://doi.org/10.5958/0976-5506.2018.01915.0>
  - [8] RAND. Indonesian Family Life Survey (IFLS). 2015. Available from: <https://www.rand.org/labor/FLS/IFLS/ifls5.html>
  - [9] RAND Corporation. The Indonesian Family Life Survey (IFLS) Santa Monica, California 2014. Available from: <http://www.rand.org/labor/FLS/IFLS.html>
  - [10] Strauss J, Witoelar F, and Sikoki B. The fifth wave of the indonesia family life survey: overview and field report. 2016, vol. 1, p. 94. Available from: [http://www.rand.org/pubs/working\\_papers/WR1143z1.html](http://www.rand.org/pubs/working_papers/WR1143z1.html)
  - [11] IPAQ. Guidelines for Data Processing and Analysis of the International Physical Activity Questionnaire (IPAQ) short and long forms, revised on November 2005. IPAQ 2005;1-15. Available from: <https://sites.google.com/site/theipaq/scoring-protocol>.
  - [12] Greenland S, Rothman KJ. Fundamentals of epidemiologic data analysis: modern epidemiology. Philadelphia: Lippincott Williams & Wilkins 2008, p. 219.
  - [13] Tsimihodimos V, Vilalpando CG, Meigs JB, Ferrannini E. Hypertension and diabetes mellitus: co-prediction and time trajectories. *Hypertension* 2018;71:422-8. <https://doi.org/10.1161/HYPERTENSIONAHA.117.10546>
  - [14] Venugopal K, Mohammed MZ. Prevalence of hypertension in type-2 diabetes mellitus. *CHRISMED Journal of Health and Research* 2014;1:223-7. <https://doi.org/10.4103/2348-3334.142981>
  - [15] de Boer IH, Bangalore S, Benetos A, Davis AM, Michos ED, Muntner P, Rossing P, Zoungas S, Bakris G. Diabetes and hypertension: a position statement by the American Diabetes Association. *Diabetes Care* 2017;40:1273-84. <https://doi.org/10.2337/dci17-0026>
  - [16] Siddiqui MA, Khan MF, Carline TE. Gender differences in living with diabetes mellitus. *Materia Socio-Medica* 2013;25:140-2. <https://doi.org/10.5455/msm.2013.25.140-142>
  - [17] Gillani SW, Ansari IA, Zaghaloul HA, Abdul MIM, Sulaiman SAS, Baig MR, Rathore H. Women with type 1 diabetes mellitus: effect of disease and psychosocial-related correlates on health-related quality of life. *J Diabetes Res* 2018; Article ID 4079087. <https://doi.org/10.1155/2018/4079087>
  - [18] Shang X, Li J, Tao Q, Li J, Li X, Zhang L, Liu X, Wang Q, Shi X, Zhao Y, Hu S, Jiang L, Yang Y. Education level, obesity and incidence of diabetes among Chinese adult men and women aged 18-59 years old: an 11-year follow up study. *Plos One* 2013;8:e66479. <https://doi.org/10.1371/journal.pone.0066479>
  - [19] Steele CJ, Schottker B, Marshall AH, Kouvonen A, O'Doherty MG, Mons U, Saum KU, Boffetta P, Trichopoulos A, Brenner H, Kee F. Education achievement and type 2 diabetes-what mediates the relationship in older adults? Data from the ESTHER study: a population-based cohort study. *BMJ Open* 2017;7:e013569. <https://doi.org/10.1136/bmjopen-2016-013569>
  - [20] Coldberg SR, Sigal RJ, Yardley JE, Riddell MC, Dunstan DW, Dempsey PC, Horton ES, Castorino K, Tate DF. Physical activity/exercise and diabetes: a position statement of the American Diabetes Association. *Diabetes Care* 2016;39:2065-79. <https://doi.org/10.2337/dc16-1728>
  - [21] Chen L, Pei JH, Kuang J, Chen HM, Chen Z, Li ZW, Yang HZ. Effect of lifestyle intervention in patients with type 2 diabetes: a meta-analysis. *Metabolism* 2015;64:338-47. <https://doi.org/10.1016/j.metabol.2014.10.018>
  - [22] Lin X, Zhang X, Guo J, Roeberts CK, McKenzie S, Wu WC,

- Liu S, Song Y. Effects of exercise training on cardio respiratory fitness and biomarkers of cardiometabolic health: a systematic review and meta-analysis of randomized controlled trials. *J Am Heart Assoc* 2015;4:4. <https://doi.org/10.1161/JAHA.115.002014>
- [23] Schellenberg ES, Dryden DM, Vandermeer B, Ha C, and Kowronyk C. Lifestyle interventions for patients with and at risk for type 2 diabetes: a systematic review and meta-analysis. *Ann Intern Med* 2013;159:543-51. <https://doi.org/10.7326/0003-4819-159-8-201310150-00007>
- [24] Yardley JE, Hay J, Abou-Setta AM, Marks SD, McGavock J. A systematic review and meta-analysis of exercise interventions in adults with type 1 diabetes. *Diabetes Res Clin Pract* 2014;106:393-400. <https://doi.org/10.1016/j.diabres.2014.09.038>
- [25] Roberts CK, Hevener AL, Barnard RJ. Metabolic syndrome and insulin resistance: underlying causes and modification by exercise training. *Compr Physiol* 2013;3:1-58. <https://doi.org/10.1002/cphy.c110062>
- [26] Magkos F, Tsekouras Y, Kavouras SA, Mittendorfer B, Sidossis LS. Improved insulin sensitivity after a single bout of exercise is curvilinearly related to exercise energy expenditure. *Clin Sci (Lond)* 2008;114:59-64. <https://doi.org/10.1042/CS20070134>
- [27] Feihl F, Liaudet L, Waeber B, Levy BI. Hypertension: a disease of the microcirculation?. *Hypertension* 2006;48:1012-7. <https://doi.org/10.1161/01.HYP.0000249510.20326.72>
- [28] Nguyen TT, Wang JJ, Islam FM, Mitchell P, Tapp RJ, Zimmet PZ, Simpson R, Shaw J, Wong TY. Retinal arteriolar narrowing predicts incidence of diabetes: the Australian Diabetes, Obesity and Lifestyle (AusDiab) Study. *Diabetes* 2008;57:536-9. <https://doi.org/10.2337/db07-1376>
- [29] Meigs JB, Hu FB, Rifai N, Manson JE. Biomarkers of endothelial dysfunction and risk of type 2 diabetes mellitus. *JAMA* 2004;291:1978-86. <https://doi.org/10.1001/jama.291.16.1978>
- [30] Ferrannini E, Cushman WC. Diabetes and hypertension: the bad companions. *Lancet* 2012;380:601-10. <https://doi.org/10.1016/S0140-6736>
- [31] Perreault L, Pan Q, Aroda VR, Barret-Connor E, Dabelea D, Dagogo-Jack S, Hamman RF, Kahn SE, Mather KJ, Knowler WC. Exploring residual risk for diabetes and microvascular disease in the Diabetes Prevention Program Outcomes Study (DPPOS). *Diabet Med* 2017;34:1747-55. <https://doi.org/10.1111/dme.13453>
- [32] Wei GS, Coady SA, Goff DC, Brancati FL, Levy D, Selvin E, Vasan RS, Fox CS. Blood pressure and the risk of developing diabetes in African Americans and whites: ARIC, CARDIA, and the Framingham Heart Study. *Diabetes Care* 2011;34:873-9. <https://doi.org/10.2337/dc10-1786>

Received on October 17, 2019. Accepted on August 24, 2020.

**Correspondence:** Aprizal Satria Hanafi, School of Public Health, University of Indonesia Depok 16424, Indonesia - E-mail: aprizalsatria-hanafi@gmail.com

**How to cite this article:** Sitorus N, Hanafi AS, Simbolon D, Martina. Joint effect of high blood pressure and physical inactive on diabetes mellitus: a population-based cross-sectional survey. *J Prev Med Hyg* 2020;61:E614-E620. <https://doi.org/10.15167/2421-4248/jp-mh2020.61.4.1406>

© Copyright by Pacini Editore Srl, Pisa, Italy

*This is an open access article distributed in accordance with the CC-BY-NC-ND (Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International) license. The article can be used by giving appropriate credit and mentioning the license, but only for non-commercial purposes and only in the original version. For further information: <https://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>*



# Joint effect of high blood pressure and physical inactive on diabetes mellitus: a population-based cross-sectional survey

## ORIGINALITY REPORT

14%

SIMILARITY INDEX

8%

INTERNET SOURCES

10%

PUBLICATIONS

11%

STUDENT PAPERS

## PRIMARY SOURCES

|   |  |    |
|---|--|----|
| 1 | Submitted to South University<br>Student Paper   | 3% |
| 2 | Anissa Rizkianti, Ika Saptarini, Rika Rachmalina. "Perceived Barriers in Accessing Health Care and the Risk of Pregnancy Complications in Indonesia", International Journal of Women's Health, 2021<br>Publication | 2% |
| 3 | Submitted to St. Petersburg College<br>Student Paper   | 1% |
| 4 | care.diabetesjournals.org<br>Internet Source   | 1% |
| 5 | Vasilis Tsimihodimos, Clicerio Gonzalez-Villalpando, James B. Meigs, Ele Ferrannini. "Hypertension and Diabetes Mellitus", Hypertension, 2018<br>Publication   | 1% |
| 6 | journals.plos.org<br>Internet Source   | 1% |

|    |   |     |
|----|---|-----|
| 7  | <a href="http://ses.library.usyd.edu.au">ses.library.usyd.edu.au</a><br>Internet Source | 1 % |
| 8  | <a href="http://www.mat.or.th">www.mat.or.th</a><br>Internet Source                     | 1 % |
| 9  | Submitted to Universitas Hasanuddin<br>Student Paper                                    | 1 % |
| 10 | <a href="http://www.nature.com">www.nature.com</a><br>Internet Source                   | 1 % |
| 11 | <a href="http://www.i-scholar.in">www.i-scholar.in</a><br>Internet Source               | 1 % |
| 12 | Submitted to Universitas Airlangga<br>Student Paper                                     | 1 % |
| 13 | Submitted to iHeed<br>Student Paper   | 1 % |

Exclude quotes  On

Exclude matches  < 1%

Exclude bibliography  On